

CONFERENCE REPORT

247

- 2 W. L. Bade and H. Jehle, An introduction to spinors, *Rev. Modern Phys.* 25:714–728 (1953).
- 3 P. Budinich and A. Trautman, An introduction to the spinorial chessboard, *J. Geom. Phys.* 4:361–390 (1987).
- 4 P. Rastall, A new spinor calculus, *Gen. Relativity Gravitation* 3:281–304 (1972).
- 5 F. A. M. Frascara and B. J. Hiley, The implicate order, algebras, and the spinor, *Found. Phys.* 10:7–31 (1980).
- 6 R. Penrose, Spinors and torsion in general relativity, *Found. Phys.* 13:325–339 (1983).
- 7 M. Riesz, *Clifford Numbers and Spinors*, Lecture Ser. 38, Univ. of Maryland, 1958.
- 8 A. Crumeyrolle, *Orthogonal and Symplectic Clifford Algebras*, Kluwer Academic, Dordrecht, 1990.
- 9 A. Crumeyrolle, The primitive idempotents on the Clifford algebras and the amorphic spinor fibre bundles, *Rep. Math. Phys.* 25:305–328 (1987).
- 10 L. D. Landau and E. M. Lifshitz, *Quantum Mechanics (Nonrelativistic Theory)*, Pergamon, Oxford, 1962.
- 11 V. Berestetski, E. M. Lifshitz, and L. Pitayevski, *Théorie Quantique Relativiste I*, MIR, Moscow, 1972.
- 12 H. Goldstein, *Classical Mechanics*, 2nd ed., Addison-Wesley, Reading, Mass., 1980.
- 13 D. Hestenes, Vectors, spinors, and complex numbers in classical and quantum physics, *Amer. J. Phys.* 39:1013–1027 (1971).
- 14 E. Schrödinger, *Space-Time Structure*, Cambridge U.P., Cambridge, 1950.
- 15 S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*, Cambridge U.P., Cambridge, 1973.
- 16 A. O. Barut, *Electrodynamics and Classical Theory of Fields*, Macmillan, New York, 1964.
- 17 A. J. Macfarlane, On the restricted Lorentz group and groups homomorphically related to it, *J. Math. Phys.* 3:1116–1129 (1962).
- 18 F. Piazzese, Spinors: An elementary point of view. Applications to the Pauli theory for the electron, to appear.

Matrices Over a Discrete Valuation Ringby ANTONIO PIZARRO¹²

The goal of this research is to establish canonical forms and similarity classes of matrices with entries in a valuation ring. As a further step, the objective is to use the similarity classes to calculate the representations of the linear group $GL(n)$ over the ring of integers in a p -field.

¹² Department of Mathematics, Centenary College, Shreveport, LA 71134.
Partially supported by the Centenary Alumni Research Award, 1990.

The case of 2×2 matrices has been solved [2]. The case of 3×3 matrices has been partially solved. Canonical forms and the similarity classes are known [3]. There are some results in the case of $n \times n$ matrices, $n > 2$.

Let R be a ring. Denote by $M_{r \times s}(R)$ the set of $r \times s$ matrices with entries in R . If R is a complete (with the p -adic topology) local Noetherian ring with maximal ideal P , then two matrices are similar over R if and only if they are similar over the Artinian rings R/P^n , $n > 1$. Canonical forms are known for matrices over R/P (since R/P is a field), namely the rational canonical form or a direct sum of rational canonical forms. The problem is to determine under what condition a matrix A over R/P will preserve the same canonical form when viewed as a matrix over R/P^n , $n > 1$. We have shown that cyclic matrices over R/P will remain cyclic over R/P^n , $n > 1$ [3].

Direct Sum of Matrices and Matrix Congruencies

IN 1952 W. E. Roth used the matrix equation $AX - XB = C$ to determine when a block upper triangular matrix in the form

$$\begin{pmatrix} A & B \\ O & C \end{pmatrix}$$

is a direct sum of matrices. The result was extended to matrices over arbitrary rings by W. H. Gustafson [1]. A similar result exists for a full matrix with entries in a complete discrete valuation ring. The proof of the following theorem has been published elsewhere [4].

THEOREM. *Let R be a complete discrete valuation ring with maximal ideal P generated by a prime p . Let $A \in M_{m \times m}(R/P^n)$ and $D \in M_{r \times r}(R/P^n)$ be fixed, and $n > 1$. For $B \in M_{m \times r}(R/P^n)$, $C \in M_{r \times m}(R/P^n)$, $B \not\equiv O \pmod{p}$, $C \not\equiv O \pmod{p}$ define*

$$\mu(B, C) = \begin{pmatrix} A & p^{n-1}B \\ p^{n-1}C & D \end{pmatrix}$$

and denote

$$\mu(O, O) = \begin{pmatrix} A & O \\ O & D \end{pmatrix}.$$

Then for $\mu(B, C)$ to be similar to $\mu(O, O) \pmod{p^n}$ it is necessary and sufficient that the congruences $B \equiv AX - XD \pmod{p}$ and $C \equiv DY - YA \pmod{p}$ have solutions.

The Similarity Classes for 3×3 Matrices

We consider the case of 3×3 matrices with entries in R/P^n , $n > 1$, modulo scalar matrices. Every matrix has a cyclic vector, i.e., it is similar to the rational canonical

forms, unless the matrix is similar to either

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \pmod{p} \quad \text{for} \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \pmod{p}$$

It has been shown [3] that in this case the matrices have a canonical form known as essentially cyclic. The form is either

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & p^k \\ a & b & c \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 0 & p^r & 0 \\ 0 & 0 & 1 \\ a & b & c \end{pmatrix}$$

for $k, r > 1$ and $a, b, c \in R/P^n$. Notice that the characteristic polynomial of the essentially cyclic matrix is congruent to $x^3 \pmod{p}$. The current stage of our research is considering $n \times n$ matrices, $n > 1$, for which the characteristic polynomial is congruent to $x^n \pmod{p}$. Our conjecture is that if the characteristic polynomial is irreducible $\pmod{p^n}$, the canonical form is essentially a cyclic matrix where the ones parallel to the main diagonal are replaced by certain powers of the prime p . In other words, the matrix is similar to

$$\begin{pmatrix} 0 & p^{k_1} & 0 & 0 & \cdots & 0 \\ 0 & 0 & p^{k_2} & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & \cdots & p^{k_{n-1}} \\ a_1 & a_2 & a_3 & a_4 & \cdots & a_n \end{pmatrix},$$

k_1, k_2, \dots nonnegative integers, $a_1, a_2, \dots \in R/P^n$. The conjecture awaits a proof or a counterexample.

REFERENCES

- 1 W. Gustafson, Roth's theorem over commutative rings, *Linear Algebra Appl.* 23:245-241 (1979).
- 2 P. C. Kutzko, On the supercuspidal representations of $GL(2)$, *Amer. J. Math.* 100(1):43-60.
- 3 A. Pizarro, Similarity of classes of 3×3 matrices over a discrete valuation ring, *Linear Algebra Appl.* 54:29-51 (1983).
- 4 ———, Block matrices over a valuation ring, *Linear and Multilinear Algebra* 21:239-241 (1987).